

# 【Instructions for the vaccination of Yellow fever】

Chubu Airport Branch Quarantine Office, Nagoya Quarantine Station

Yellow fever is an acute viral hemorrhagic disease **transmitted by mosquito**.

The disease is endemic in tropical areas of Africa and Central and South America.

It is preventable by **vaccination and personal preventive measure against mosquito bite**.

It is recommended to get vaccination, if you travel to areas at risk of yellow fever.

In some countries or areas, it is required to show according to each country's regulation.

## ● Yellow fever vaccine

Yellow fever vaccine is made from the virus of yellow fever.

The virus is weakened not to cause real yellow fever (**attenuated live vaccine**).

The doctor in charge of vaccination gives yellow fever vaccine a subcutaneous injection.

Preventing effect (immunity) is now thought to continue for lifelong by single shot of the vaccine.

## ● Side effect of yellow fever vaccine

### ➤ Mild side effects

Some of the following symptoms may occur within a few days after vaccination in approximately 10% of vaccinees.

**Redness, swelling or pain at the injected site /  
Headache / Fever / Muscle pain / Back pain / Joint pain / Fatigue / Rash**

These side effects are within normal reactions when your body gets immunity against yellow fever.

These symptoms will usually disappear within a few days spontaneously.

If these symptoms continue and/or affect your daily life, seek medical advice from your doctor, as those symptoms might be due to other diseases.

### ➤ Severe side effects

#### • Anaphylactic shock

Symptoms are rash, dyspnea, vomiting, sudden drop of hypotension and so on.

It may **rarely (1 in 100,000)** occur **within 30 minutes after vaccination**.

Therefore, vaccinee is supposed to stay in the waiting room for 30 minutes after vaccination.

#### • Disorder of brain and nerves / Disorder of internal organs

Severe side effects may occur within 1 month after vaccination in a few cases in 100,000 vaccinees.

If high fever over 38.0 degrees Celsius (100.4 in Fahrenheit) continues or seizure/convulsion occurs, seek medical advice immediately as well as let us know by phone (0569-38-8205).

《For women》

### ➤ **Avoid pregnancy for 2 months** after vaccination.

### ➤ **Avoid breast-feeding** after vaccination unless you and your breast-fed child get this vaccination simultaneously. If you have any questions, consult the doctor in charge of vaccination.

**To guardian**

Read "Instructions for the vaccination of Yellow fever" before filling out this form.

## Guardian's Agreement Form

To the Chief of Chubu Airport Branch Quarantine Office

I have read "Instructions for the vaccination of Yellow fever" and fully understood its benefits and side effects.

I agree that the vaccinee (**Name** \_\_\_\_\_) gets vaccine of yellow fever on (**(Year)20** \_\_\_\_\_ **(Month)** \_\_\_\_\_ **(Day)** \_\_\_\_\_).

\_\_\_\_\_ **(Year)20** \_\_\_\_\_ **(Month)** \_\_\_\_\_ **(Day)** \_\_\_\_\_

**【Guardian】**

Address \_\_\_\_\_

Phone Number \_\_\_\_\_

Name \_\_\_\_\_

Guardian's  
relationship to  
vaccinee

**【Vaccinee】**

Address \_\_\_\_\_

Name \_\_\_\_\_

**To guardian**

Read "Instructions for the vaccination of Yellow fever" before filling out this form.

**Guardian's Agreement Form**

**Example**

To the Chief of Chubu Airport Branch Quarantine Office

I have read "Instructions for the vaccination of Yellow fever" and fully understood its benefits and side effects.

I agree that the vaccinee (**Name** \_\_\_\_\_) gets vaccine of yellow fever on (**(Year)20** \_\_\_\_\_ **(Month)** \_\_\_\_\_ **(Day)** \_\_\_\_\_).

**(Year) 20** \_\_\_\_\_ **(Month)** \_\_\_\_\_ **(Day)** \_\_\_\_\_

【Guardian】

Fill in "Date of vaccination"

Address \_\_\_\_\_

Phone Number \_\_\_\_\_

Name \_\_\_\_\_

Guardian's relationship to vaccinee

Signature of guardian

Fill in phone number we can get connection on date of vaccination

【Vaccinee】

Address \_\_\_\_\_

Name \_\_\_\_\_

Signature of vaccinee